أشعة تصوير الرحم بالصبغة

Hysterosalpingogram





مستشفيات السلام AL SALAM HOSPITALS TRUST تحت امانة
1830003
alsalamhospt
alsalamhosp.com

Hysterosalpingogram "HSG" is a radiological / medical procedure that can be resorted to in certain cases, including uterus problems. Check the following details:

What is a Hysterosalpingogram?

A Hysterosalpingogram (HSG) is an X-ray procedure where contrast is used to look at the inside of your fallopian tubes and uterus. The contrast allows your uterus and fallopian tubes to be seen more clearly on the x-ray pictures.

Will there be any discomfort and is any anesthetic needed?

Mild discomfort may occur when the catheter is inserted into your cervix. Mild pain may occur when the contrast is injected and flows through your fallopian tubes.
If you need the procedure to be done with anesthesia, kindly refer to a Customer Relation staff member in the Reception Area to guide you with the appointment procedure.

Consent

A Radiologist will explain the procedure to you and you will be asked to sign a consent form for the Hysterosalpingogram examination.

Preparation for the procedure

To have this procedure and for your own safety you must:

- Talk to your referring doctor if you have an active sexually transmitted disease or a pelvic infection.
- NOT to be pregnant. You may be asked to take a pregnancy test prior to the procedure.
- NOT be on your period. The procedure is usually done between the end of your menstrual flow and day 12 of your cycle. Please note the start date of your last menstrual period.

أشعة تصوير الرحم بالصبغه هـي نـوع مـن الفحوصـات والإجـراءات الطبيـة التـي قـد يتـم اللجـوء إليهـا في حـالات معينـة مـن ضمنهـا مشـاكل الرحـم، إليك كافة التفاصيل بشأنها :

ما هو فحص تصوير الرحم بالصبغة (HSG)؟

هـو فحـص تصويـر الرحـم و الأنبـوب الفالـوبي بواسـطة الأشـعة السـينية و المادة الصبغيـة حيـث أن المادة الصبغيـة تسـمح برؤيـة الرحـم و الأنبـوب الفالـوبي بوضـوح ودقة.

هل هناك شعور بعدم الراحة؟ هل يحتاج الفحص للتخدير؟

سيكون هناك إحساس بعدم الراحة حين إدخال الأنبوب (catheter) داخل عنق الرحم، و من الممكن الشعور بألم بسيط حين يتم حقن المادة الصبغية وتدفقها خلال الأنبوب الفالوبي.
إذا كنتي تحتاجين للتخدير يرجى طلب ذلك من موظفي خدمة العملاء بالإستقبال لتنسيق حجز المواعيد.

إقرار

سيقوم طبيب الأشعة بشرح الفحص لك ثم سيطلب منك التوقيع على نموذج الموافقة على الفحص.

التجهيز للفحص:

لإجراء الفحص ولسلامتك، يرجى منك التالي:

- التحدث مع طبيبك المعالج في حال تعانين من مرض منقول جنسيا أو إلتهاب بالحوض.
- أن لا تكوني حامل، وقد تحتاجين لعمل تحليل الحمل قبل الفحص.
- أن لا يكون الفحص خلال الدورة الشهرية (غالبا يتم عمل الفحص بالأيام بين إنتهاء التدفق و اليوم ال 12 من الدورة).

During the procedure:

You will lie down on a table, in the same position as a pelvic examination.
A speculum is placed inside your vagina so that your cervix can be visualized. An instrument is attached to the cervix and then a cannula is placed in the outer opening of the cervix.
Through this tube a contrast is injected into the uterine cavity.
X-ray pictures are taken and reviewed.
The speculum and the cannula are removed at the end of the procedure.

After the procedure

You may have 'spot bleeding' and/ or slight watery discharge and will need to wear a sanitary pad (not a tampon).
You may have mild abdominal pain for 1 to 2 days after the procedure. Pain reliever medications may be taken to relieve mild abdominal pain.

Common risks and complications include:

- Bleeding from the vagina.
- Dizziness or feeling faint, this is usually resolved with bed rest.

Less common risks and complications include:

- Infection, requiring antibiotics and further treatment.
- Damage to the fallopian tubes, requiring corrective surgery.

Rare risks and complications include:

- Allergic reaction to the contrast. This could result in a rash, hives, itching, nausea, fainting or shortness of breath. Medication may be given to relieve this.

What are the safety issues when you leave the hospital?

Go to the nearest Emergency Department if you become unwell or have:

- Severe or increased pelvic pain.
- Heavy vaginal bleeding (more than a pad an hour).
- Vomiting.
- Fever.
- Foul smelling or odd vaginal discharge.

خلال الفحص :

تستلقين على سرير الفحص، كوضعية فحص الحوض. يتم إدخال منظار داخل المهبل لرؤية عنق الرحم و حقن المادة الصبغية إلى تجويف الرحم من خلاله. يتم أخذ صور بالأشعة السينية وعند الإنتهاء من التصوير، يتم إخراج المنظار والأنبوب.

بعد الفحص :

قد يوجد بقع دم مع فقدان السوائل بسبب المادة الصبغية في الرحم، مما يستلزم وضع فوط صحية. وقد تشعري بألم بسيط في البطن لمدة يوم أو إثنين، حيث يمكنك أخذ المسكنات لها .

ما هي مخاطر هذا الفحص بالتحديد؟

مخاطر ومضاعفات شائعة للفحص تتضمن:

- نزيف المهبل
- دوخة أو إغماء. قد يختفي مع الاستلقاء والراحه ع السرير.

مخاطر ومضاعفات غير شائعة للفحص:

- التهاب يحتاج لمضاد لعلاجه
- ضرر بالأنبوب الفالوبي (يحتاج لعملية تصحيحية).

مضاعفات نادرة:

حساسية من المادة الصبغية قد تنتج عنها طفح جلدي، قشعريرة، حكة، غثيان، إغماء، ضيق بالتنفس. سيتم إعطاء أدويه لعلاجه.

حرصا لسلامتك بعد الخروج من المستشفى:

زوري أقرب قسم طوارئ إذا ظهر عليك هذة الأعراض:

- ألم حاد بالحوض.
- نزيف كثير بالمهبل (يستلزم تغيير الفوطة الصحية كل ساعة).
- الإستفراغ.
- سخونة (ارتفاع الحرارة).
- رائحة كريهة أو إفرازات مهبلية غريبة.